

Keeping Hydrated

You should aim to have at least 1.6 - 2 litres (around 6 - 8 glasses) of fluid per day to stay hydrated. Keeping hydrated can prevent or aid the treatment of constipation, low blood pressure, urinary tract infections (UTIs), pressure ulcers and falls.



Did you know? Around 20% of our daily fluid intake comes from within our food. See *'Hydration Boosters'* resource for more information.

All fluids count, except for alcohol! Choose a drink that you are most likely to enjoy and finish.

Do not wait until you feel thirsty to have a drink; thirst is a late response to dehydration.

Use the pee chart to score your hydration status as the colour of your urine can indicate dehydration risk.

Serve drinks at their optimum temperature and replenish any drink that has been left to stand.

Choose nourishing fruit or milk based drinks if you are not eating well or need to gain weight (see 'Super Shakes' resource for recipe examples).

Opt for water, skimmed milk or sugar-free drinks if you have diabetes or are trying to lose weight.

Replace fluid that is lost through sweat, open wounds, diarrhoea or vomit to prevent dehydration.

You will sweat more in warm conditions and when you are more active.

Limiting your fluid intake can make incontinence worse because it reduces your bladder's capacity.

Increase your fluid intake earlier in the day if you worry about urinating at night.

Fit your fluid intake around your daily routine and take a bottle with you on the go.

Spout cups, handled mugs or plastic tumblers may be lighter and easier to handle.

Healthy pee is 1-3 4-8 you must hydrate!		
	1	

2

3

4

5

6

7

8



Tip: Use a measuring jug to gauge the volume of your cups/glasses at home and complete a 'Self-Reported Fluid Intake' resource to record your daily fluid intake.